

# Rescon Tablets

NDC 64543-097-90

## ***Drug Facts***

### ***Active Ingredients (per tablet)***

Dexchlorpheniramine Maleate 2 mg

Pseudoephedrine Hydrochloride 60 mg

### ***Purpose***

Dexchlorpheniramine Maleate Antihistamine

Pseudoephedrine Hydrochloride Nasal Decongestant

### ***Uses***

For the temporary relief of

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes due to hay fever or other upper respiratory allergies
- nasal congestion

Temporarily helps

- clear nasal passages
- shrink swollen membranes

### ***Warnings***

- **Do not exceed recommended dosage**
- **May cause excitability especially in children**
- May cause drowsiness; alcohol, sedatives, and tranquilizers may increase the drowsiness effect.

### ***When using this product***

- Avoid alcoholic beverages
- Use caution when driving a motor vehicle or operating machinery.

### ***Do not use***

- If you have a breathing problem such as emphysema or chronic bronchitis
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

**Stop use and ask a doctor**

- If nervousness, dizziness, or sleeplessness occur.
- If symptoms do not improve within 7 days or are accompanied by fever.

**Ask a doctor before use if you have**

- glaucoma
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- difficulty in urination due to enlargement of the prostate gland
- or if you are taking sedatives or tranquilizers

**If pregnant or breast feeding**

Ask a health professional before use.

**Keep out of the reach of children**

In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

***Directions***

- **Adults (12 and older):** One tablet every 4 to 6 hours. Not to exceed 4 doses in 24 hours.
- **Children under 12 years of age:** Consult a physician.

***Other information***

store at 20° -25° C (68° -77° F)

***Inactive ingredient***

colloidal silicon dioxide, crospovidone polyplasdone, D&C Red #30, D&C Yellow #10, FD&C Blue #1, lactose monohydrate, magnesium stearate, microcrystalline cellulose, and povidone